

ROLE OF BIOSTATISTICS IN CLINICAL TRIALS FOR NUTRITIONAL INTERVENTION

Iqra Zafar^{1*}, Sidratulmuntaha¹, Laiba Qasim¹, Hajra Mazher¹, Laiba Mueen¹

¹Department of Human Nutrition and Dietetics (HND), The Islamia University of Bahawalpur (IUB), Pakistan

*Corresponding author e-mail: iqra.zafar@gmail.com

ABSTRACT In this article, we will discuss the applications of biostatistics in clinical trials and nutritional intervention. Its role in modeling nutrient effects on health outcomes, randomized controlled trials in nutrition research, and investigating nutrient gene interaction. Biostatistics contributes to the design of clinical trials and analysis of data. Biostatistics also tells us about the relationship between dietary patterns and health outcomes. It undergoes meta-analysis of nutrition intervention studies. It also investigates nutrient-environment interaction. It also evaluates the cost-effectiveness of nutritional intervention. The role of biostatistics in nutritional intervention is emerging day by day. It provides us with evidence-based contributions in clinical trials and nutritional intervention.

Keywords: *Biostatistics; Trials; Nutrition; Randomized; Interventions*

INTRODUCTION To promote human health, nutrition is essential. Many non-communicable diseases can today be prevented and treated in part due to the influence of nutrition. The quality of the evidence supporting nutritional guidelines is in doubt because most nutrition evidence is derived from observational study results, even though nutrition plays a significant role in health and the amount of information in this area has grown recently. Furthermore, due to design limitations, the results of randomized clinical trials (RCTs) based on nutritional therapies are sometimes disputed or have little therapeutic importance. A series of steps intended to modify a population's or an individual's nutritional status is known as a nutritional intervention. A nutritional education program can be implemented or nutrients can be administered, among other things, as part of a nutritional intervention. There are four categories of dietary interventions according to the World Health Organization (WHO):

1. Behavioral therapies seek to alter eating patterns through behavioral means.
2. Nutrition is added to basic meals through a process known as fortification.
3. Supplementation is the process of giving a certain nutrient to a particular group of people.
4. Regulatory interventions aim to adjust nutrition and promote health by regulating specific behaviors.(Martínez-López, Pérez-Guerrero et al. 2022)

In addition to the research on specific nutrients, there needs to be research on the health impacts of dietary patterns. Dietary patterns take into account the intricate links that exist between various foods or nutrients in their entirety, mirror the dietary practices of individuals, and offer additional insight into the instances in which multiple nutrients are linked to specific diseases. Dietary pattern analysis is therefore viewed as an adjunctive tool to the investigation of individual nutrients. Statistical techniques that fully utilize dietary data gathered from various populations to establish dietary trends have been developed in the last several decades.(Zhao, Li et al. 2021)

Biostatistics Application in Clinical Trials

Clinical trial design, execution, and reporting all heavily rely on biostatistics. The understanding and application of biostatistics have significantly increased in the last few years. In addition to being in charge of the quantitative aspects of clinical trial designs and the analysis of trial data, biostatisticians are esteemed members of the clinical teams who participate in strategic conversations and assist in resolving day-to-day problems with studies and clinical programs. The innovative statistical approach is developed and implemented mostly by biostatisticians (Marchenko, LaVange et al. 2020) The majority of statistical techniques for clinical trial design and analysis were created in the context of confirmatory trials with sample sizes that were comparatively large—possibly hundreds or even thousands of patients.(Hilgers, Roes et al. 2016) processed, deep-fried foods, and beverages with added sugar was linked to a greater chance of obesity. Nevertheless,

considering the variations in eating customs and cultures(Jiang et al., 2022).

Modeling Nutrient Effects on Health Outcomes

The health care system is heavily burdened financially by excess body weight, high blood pressure, hyperlipidemia, impaired glucose tolerance, and the chronic diseases they are associated with. Dietary modifications can alter these risk factors, but long-term sustainability and success in dietary intervention studies have generally been poor. Dietary intervention studies with sufficient sample size, duration, and acceptable compliance have been proven to be economically feasible. Furthermore, national dietary recommendations are largely ignored by diets around the world.(Grieger, Johnson et al. 2017) Nutrition epidemiologists can acquire data encompassing extensive biological, psychological, behavioral, and demographic characteristics of public health importance through national health surveys.(Hartwell, Khojasteh et al. 2019).

Randomized Controlled Trials in Nutrition Research

Dietary recommendations that are supported by evidence should be founded on the entirety of the available data. In nutrition research, systematic reviews and meta-analyses are the most beneficial, trustworthy, and impartial methods for compiling data related to a particular study issue. The lack of evidence-quality evaluation in numerous meta-analyses has reduced our confidence in the apparent effect.(Schwingshackl, Knüppel et al. 2016) Randomized controlled trials (RCTs) are the gold standard in the field of human nutrition when it comes to proving a causal relationship between exposure to foods, nutrients, or dietary patterns and predetermined outcome measures like body composition, biomarkers, or event rates. (Lichtenstein, Petersen et al. 2021).

Investigating Nutrient Gene Interaction

The long-term causal effects of different exposures on clinical and epidemiological outcomes are estimated using observational data and the Mendelian randomization technique, which is widely used. To eliminate bias brought on by confounding and reverse causality, it makes use of genetic variants.(Burgess, Daniel et al. 2015).

Analysis of Dietary Patterns and Health Relationship

The association between food insecurity, eating patterns, and weight increase is not well known, but it may be linked to both unhealthy weight and bad diet. Developing and putting into practice successful behavioral and lifestyle treatments for at-risk populations may benefit greatly from a deeper comprehension of this link. Food insecurity has been associated in the past with poor diet quality and reduced consumption of healthy food groups, especially when it comes to the amount of fruits and vegetables consumed.(Morales and Berkowitz 2016) There is little information on the relationships between food insecurity, economic disadvantage, and abnormal weight (underweight, overweight, or obese). Malnutrition, on the other hand, can be brought on by a persistent lack of financial resources and food instability. Biostatistics has indicated a correlation between food insecurity and several variables, such as residing in a low-income neighborhood, having a low income, and experiencing unemployment.(Moradi, Mirzababaei et al. 2019).

Meta-Analysis of Nutrition Intervention Studies

Obesity is acknowledged as a global health concern. Public health interventions over the past three decades have not only failed to stop the growth in population weight but also to limit its rate of increase. Dietitian professional associations, like the USA's Academy of Nutrition and Dietetics, offer guidelines to their members for patient management of obesity. Authorities at the national and international levels, including the World Health Organization, have assessed the data about successful adult overweight and obesity interventions and have established target intervention outcomes. Remarkably, and perhaps concerningly, given the relative availability of evidence-based guidelines and systematic reviews on managing obesity, including those issued by dietetic associations. (Williams, Barnes et al. 2019).

Investigating Nutrient environment interaction

Gaining more understanding of the genetic diversity and G-H-E interaction in foliar nutritional characteristics and growth traits is also important (as well as their interactions). Examining potential applications of physiological markers to tree-selection programs is the initial step.(Li, Xue et al. 2015) . Another example is that of soil and nutrient interaction. The effects of adding limiting resources, such as water and nutrients, on the growth and Eco- - physiological processes in southern USA pine plantations have been the subject of much investigation. The more the nutrients, the more plant growth will be. (Will, Fox et al. 2015).

Evaluating the Cost-effectiveness of Nutritional Intervention

To summarize all the data and determine whether an intervention is worth the money invested in it, the cost-effectiveness plane is given. With the limited resources at hand, the process of economic evaluation can make a significant contribution to the development and identification of cost-effective treatments, assisting in the pursuit of the greatest possible reduction in malnutrition. When designing and assessing dietary interventions, the health economics field's cost-effectiveness framework can be a valuable resource. It compels the assessors to investigate the numerous factors affecting the intervention's efficacy in a practical setting. It also considers the various financial factors that have an impact on an individual's health and well-being.(Wieser and Tzogiou 2017).

CONCLUSION

In conclusion, the field of nutrition research presents a complex landscape where various factors interact to influence human health. Despite challenges such as evidence quality and study design limitations, randomized controlled trials stand out as indispensable tools for establishing direct causal relationships between nutrition and health outcomes. However, it's crucial to recognize that nutrition extends beyond the mere consumption of individual nutrients; understanding dietary patterns and their intricate associations with health outcomes calls for a holistic approach to nutrition science. Moving forward, advancements in biostatistics, coupled with interdisciplinary collaboration, will continue to drive progress in nutrition science. By leveraging innovative statistical methods and integrating insights from

genetics, epidemiology, and behavioral science, we can develop more nuanced and tailored approaches to nutrition interventions.

REFERENCES

- Martínez-López, E., et al. (2022). "Methodological aspects in randomized clinical trials of nutritional interventions." *Nutrients* 14(12): 2365
- Zhao, J., et al. (2021). "A review of statistical methods for dietary pattern analysis." *Nutrition Journal* 20: 1-18.
- Marchenko, O. V., et al. (2020). "Biostatistics in Clinical Trials." *Quantitative Methods in Pharmaceutical Research and Development: Concepts and Applications*: 1-70.
- Burgess, S., et al. (2015). "Network Mendelian randomization: using genetic variants as instrumental variables to investigate mediation in causal pathways." *International journal of epidemiology* 44(2): 484-495.
- Grieger, J. A., et al. (2017). "Evaluation of simulation models that estimate the effect of dietary strategies on nutritional intake: a systematic review." *The Journal of nutrition* 147(5): 908-931.
- Hartwell, M. L., et al. (2019). "Using structural equation modeling to examine the influence of social, behavioral, and nutritional variables on health outcomes based on NHANES data: addressing complex design, nonnormally distributed variables, and missing information." *Current developments in nutrition* 3(5): nzz010.
- Hilgers, R.-D., et al. (2016). "Directions for new developments on statistical design and analysis of small population group trials." *Orphanet journal of rare diseases* 11: 1-10.
- Li, Y., et al. (2015). "Genetic parameters and clone by environment interactions for growth and foliar nutrient concentrations in radiata pine on 14 widely diverse New Zealand sites." *Tree Genetics & Genomes* 11: 1-16.
- Lichtenstein, A. H., et al. (2021). "Perspective: design and conduct of human nutrition randomized controlled trials." *Advances in Nutrition* 12(1): 4-20.
- Moradi, S., et al. (2019). "Food insecurity and adult weight abnormality risk: a systematic review and meta-analysis." *European journal of nutrition* 58: 45-61.
- Morales, M. E. and S. A. Berkowitz (2016). "The relationship between food insecurity, dietary patterns, and obesity." *Current nutrition reports* 5: 54-60.
- Schwingshackl, L., et al. (2016). "Perspective: NutriGrade: a scoring system to assess and judge the meta-evidence of randomized controlled trials and cohort studies in nutrition research." *Advances in Nutrition* 7(6): 994-1004.
- Wieser, S. and C. Tzogiou (2017). "Does it work? Is it worth it? Evaluating the costs and benefits of nutritional interventions." *Sustainable Nutrition in a Changing World*: 343-353.
- Will, R. E., et al. (2015). "A range-wide experiment to investigate nutrient and soil moisture interactions in loblolly pine plantations." *Forests* 6(6): 2014-2028.
- Williams, L. T., et al. (2019). "How effective are dietitians in weight management? A systematic review and meta-analysis of randomized controlled trials." *Healthcare*, MDPI.